

Xpert Hotstart Mastermix (2X)
#GE18.0100 (1.25 ml) | #GE18.5100 (6.25ml)
(for research only)

Product:	Xpert Hotstart DNA polymerase is a chemically modified hotstart Taq DNA polymerase, with excellent amplification efficiency, enabling higher specificity, increased sensitivity, and greater yield, as compared to standard Taq DNA polymerases, making this the ideal choice for consistent results in complex PCR amplifications and multiplex PCR.
Source:	Recombinant, purified from <i>E.coli</i>
Quantity:	Xpert Hotstart DNA polymerase is supplied as a convenient 2x mastermix, containing all required components for PCR, except specific primers. Final concentration of MgCl ₂ will be 1.5mM. Each vial is supplied with an additional separated 25mM MgCl ₂ allowing further optimization. #GE18.0100 contains 1.25 ml of Xpert Hotstart Mastermix (2X) #GE18.0500 contains 5x 1.25ml of Xpert Hotstart Mastermix (2X)
Applications:	Complex targets (including GC-rich and AT-rich templates), Multiplex PCR.
Properties:	Amplicon size: up to 5kb Extension Rate 2kb/min Hotstart: Yes A-overhang: Yes
QC:	Functionally tested in PCR. Absence of endonucleases, exonucleases, and ribonucleases was confirmed by appropriate assays.
Storage:	-20°C for at least 1 year. No loss of performance is detected after 20 freeze/thaw cycles.

GRiSP Research Solutions
Rua Alfredo Allen, 455
4200-135 Porto
Portugal
www.grisp.pt | info@grisp.pt

Basic Protocol


Optimal PCR cycling conditions (incubation times and temperatures) depend on DNA target (GC-content, size, quantity, purity, etc) and specific primers and need to be determined case by case. Xpert Hotstart Mastermix (2X) includes dNTPs, MgCl₂ and other components to maximize success rates. We suggest to start with the following basic protocol and subsequently, if desired, optimize annealing temperature, Mg²⁺ concentration, incubation times and cycling number.

Mix for each PCR reaction according to the following table. In order to minimize risk of contamination, reagent loss and improve pipetting accuracy, we recommend to prepare a mastermix for multiple samples (N), always including a negative control for the detection of possible contaminants, by mixing all components (N+1), except template DNA, dividing the mixture equally into each tube and then add template DNA or PCR grade water in case of the control to the individual PCR tubes. For smaller/larger reaction volumes, scale it down/up proportionally.

Component	Volume (25µl)	Final Conc.
Xpert Hotstart Mastermix (2X)	12.5 µl	1X
Forward primer (5 pmol/ µl)	2 µl	0.4 µM
Reverse primer (5 pmol/ µl)	2 µl	0.4 µM
Template DNA	0.25 - 10 µl	1-250 ng*
PCR –grade water	up to 25 µl	

*) In case of cDNA <50ng and in case of gDNA <250ng (total amount).
For smaller/larger reaction volumes, scale it down/up proportionally.

Set-up initial PCR amplification as follows:



After an initial cycle of 15 min at 95°C (cleavage of the chemical moiety), cycle 25-35 times for 15-30 seconds at 95°C, 15-30 seconds at Ta, and 30 seconds per kb of target DNA at 72°C for extension. Set the annealing temperature (Ta) as the melting temperature (T_m) of the primer with the lowest T_m. Include a final extension step of 5 min at 72°C to ensure that all amplicons are fully extended and include 3'-A-overhang. Analyze PCR products by DNA Agarose gel electrophoresis.

Optimization

Annealing Temperature (Ta) and Primers

Optimizing the annealing temperature is crucial, especially in case of multiplex PCR, as a too low temperature might result in non-specific amplification whereas a too high temperature results in no amplification. The melting temperature (T_m) is defined as the temperature in which 50% of the primer and its complementary sequence of the target DNA are present as duplex DNA. By increasing the temperature above the melting temperature, this percentage decreases, however, primers will still anneal (up to a certain point) and initiate extension. PCR can therefore be performed at temperatures of several degrees higher than T_m and it is therefore recommended to optimize the Ta by performing a temperature gradient (e.g. starting at the lowest T_m or a few degrees below and increasing with 2°C increments). Ideally, primers have melting temperatures of approximately 60°C and final concentration should be between 0.2 and 0.6µM (each).

MgCl₂

Xpert Hotstart Mastermix (2X) contains 3mM MgCl₂ (final concentration is 1,5mM). This product is supplied with a separated vial of 25mM MgCl₂ allowing for the optimization of the final MgCl₂ concentration, which is normally between 1,5mM and 3mM

Incubation times and number of cycles.

Denaturation and annealing steps may require less time depending on the thermocycler apparatus (ramp rate), reaction volume and PCR tube (varies with the efficiency of heat-transfer). It might be worthwhile to optimize (reduce) times to as low as 10 seconds for both denaturation and annealing steps, which will greatly reduce overall PCR time. Faster cycling conditions should not be applied in multiplex PCR.